

Spectroscopic Window on Tumor Metabolism

Michael Garwood, Ph.D.

Patrick Bolan, Ph.D.

Greg Metzger, Ph.D.

Center for Magnetic Resonance Research, Department of Radiology, and Cancer Center,
University of Minnesota Medical School, Minneapolis, Minnesota 55455 USA

Introduction

In vivo MRS can be performed along with MRI to measure metabolic properties of tumors. MRS has proven its value as a research tool for elucidating metabolic and biochemical aspects of cancer. In addition, *in vivo* MRS holds much promise for becoming an indispensable clinical tool for detecting, diagnosing, and monitoring responses to therapies in patients. Well known examples of the latter include the use of MRS to improve the accuracy in differentiating between malignant and benign lesions and to detect an early response to a chosen therapy. At present, a growing number of radiologists as well as oncologists are interested in incorporating MRS into their MRI protocols. This article introduces ^1H MRS of cancer, discusses current methods and technical issues, and describes some of the more common applications to cancer. Although *in vivo* MRS has been used to study cancers in many different parts of the body, the greatest progress in the development and validation of MRS has occurred mainly for three types of cancer: brain, breast, and prostate. Thus, the present discussion will focus mainly on these cancers, although promising results have been obtained in pilot ^1H MRS studies of cancers in other sites, including extracranial lymphoma and germ cell tumors [1], hepatic tumors [2], head and neck tumors [3], melanoma, and lymph nodes [4].

Altered ^1H metabolites in cancer

Malignant tissues have unique spectral features that distinguish them from normal tissues. Many of the metabolite resonances present in the normal host tissues are absent or decreased in malignant lesions. For example, resonances from polyamines and citrate (Cit), which are hallmarks of normal healthy and benign hyperplastic prostatic tissues, are absent or substantially reduced in prostate cancers. Likewise, resonances from N-acetylaspartate (NAA), a neuronal marker, are typically reduced in brain tumors. Total creatine (tCr), an indicator of bioenergetic status, is another resonance that typically has altered intensity in cancers. Other hallmarks of cancer include increased levels of choline-containing compounds (tCho), mobile lipids, and lactic acid. Due to their importance in cancer, each of these compounds is discussed in further detail below.

Choline compounds

Numerous *in vivo*, *ex vivo*, and *in vitro* studies have demonstrated elevated levels of tCho in neoplastic tissues and cells. *In vivo* ^1H spectra of cancerous lesions typically display a

prominent resonance from choline compounds at 3.2 ppm. *Ex vivo* studies have been performed to identify the different choline compounds giving rise to this resonance. In breast cancer, high-resolution ^1H spectra acquired from biopsy tissues have shown that the 3.2 ppm resonance is actually a superposition of several resonances [5-7]. The primary constituents are those with a trimethylamine moiety $[\text{R}-(\text{CH}_2)_2-\text{N}^+(\text{CH}_3)_3]$, including free choline (Cho), phosphocholine (PCho), and glycerophosphocholine (GPC). Other metabolites possibly contributing include taurine, glucose, phosphoethanolamine, and myo-inositol (mIns) [7]. The choline head groups associated with semi-mobile lipids may also contribute. These resonances can be separated in *ex vivo* studies with high-resolution NMR spectrometers, but *in vivo*, these peaks are substantially broadened, and at fields as high as 4 Tesla these resonances are generally indistinguishable. Consequently, the simplified approach used in *in vivo* studies is to treat the 3.2 ppm spectral peak as a single resonance, and thus it is referred to as tCho (total choline-containing compounds).

The precise mechanisms that produce elevated tCho levels in cancers are as yet not fully understood. A working hypothesis is that elevated tCho is an indicator of increased cellular proliferation [8, 9]. Indeed, the largest component contributing to the tCho peak from neoplastic tissue is usually PCho, a known precursor of membranes. Another major component, GPC, is a breakdown product of membranes. It is known that tCho levels can be modulated by numerous changes in enzymatic activity and fluxes in these anabolic and catabolic pathways [10, 11], and the increased tCho level in neoplastic tissues may reflect increased membrane turnover in neoplastic tissues. In the author's opinion, the present understanding of elevated tCho in malignant cells remains an oversimplified view, and further research is needed to obtain a complete picture of the biological processes leading to elevated tCho. It can be expected that MRS will likely play a vital role in elucidating these processes.

Lipids

Resonances from mobile lipids can be observed in ^1H spectra of breast [12], brain [13], and prostate tumors [14]. The amplitude of the lipid resonance can vary dramatically depending on the tissue heterogeneity. Lipid peaks in tumor spectra can arise from sources other than normal adipose tissue. For example, resonances from mobile lipids have been shown to occur in necrotic tissues [15]. Furthermore, experimental data from *in vitro* cell studies suggest the formation of intracellular lipid droplets are a source of lipid resonances [16]. In studies of T-cell lymphoblast cultures, a correlation between the methylene-to-methyl signal ratio and number of apoptotic cells was found [17]. In accordance with this finding, lipid levels were found to correlate with apoptosis and early cell death in rat BT4C gliomas [18]. The presence of elevated lipid resonances in human astrocytic tumors has been suggested to have prognostic significance [19].

In human breast, the adipose tissue which is not directly involved in the carcinoma can pose significant problems for *in vivo* ^1H MRS. When trying to choose the volume-of-interest (VOI) for localized MRS studies of breast cancer, any adipose tissue inadvertently included in the VOI creates a partial volume effect, reducing the effective volume for spectroscopy. Likewise, in studies of prostate and brain tumors, it is very important to suppress intense signals from adipose tissues which surround the prostate and brain, in order to avoid spectral contamination ("bleed effect") and artifacts (e.g., baseline distortion). Adipose tissue also limits the ability to optimize (or "shim") the homogeneity of the magnetic field inside the VOI, which in turn leads to broad resonances and reduced signal-to-noise ratio. Intense lipid resonances can also produce sideband artifacts which can interfere with MRS measurements [20]. These artifactual resonances can be larger than the tCho resonance. To reduce sideband artifacts, our

group uses a method called echo-time (TE) averaging, which causes coherent cancellation of sideband artifacts by averaging spectra acquired at several different TE values [20].

Lactate

Under hypoxic conditions, neoplastic cells are thought to derive energy primarily from glycolysis. Even in the presence of oxygen, malignant lesions can have an increased glycolytic metabolism. Many investigations of experimental tumors have provided evidence of reduced respiratory capacity and increased reliance on glycolysis for energy production. Consequently, the concentration of lactate in tumors is generally higher than that in normal tissues. In single-voxel ^1H MRS measurements on normal human brain the lactate concentration was ~ 0.6 mmol/kg [21], whereas in untreated human brain tumors the lactate levels were found to be 2.6 ± 0.8 mmol/kg (mean \pm S.D., $n=7$) [22]. In studies of rat C6 glioma, we observed a positive correlation between the tumor lactate levels measured by *in vivo* ^1H MRS and the neoplastic cell density determined by histopathology [23]. Of interest, *ex vivo* studies (non-MRS) have revealed a positive correlation between the incidence of metastasis and the mean lactate concentration in biopsy specimens of cervical and head and neck tumors [24, 25].

Applications

Breast Cancer

The first *in vivo* MRS studies of breast measured resonances from ^{31}P nuclei. These studies showed that measurable variations in phospholipid metabolism could be detected and used for diagnosing cancer and monitoring response to treatment (reviewed in [26, 27]). More recently, there has been growing interest in breast cancer research using ^1H MRS, due to its higher sensitivity than ^{31}P MRS. The first breast ^1H MRS reports focused on the diagnostic utility of the water:fat ratio in the breast [28-30], but subsequent studies did not find this ratio to be a useful diagnostic metric [31, 32]. However, a number of studies performed with ^1H MRS noted the presence of tCho in spectra of malignant lesions, but not in benign or normal tissues [29, 31-36].

The majority of breast MRS studies to date have used single-voxel spectroscopy (SVS) to localize the spectrum to a single volume centered on the lesion of interest. While most breast MRS studies have been done with SVS, other researchers have explored the use of spectroscopic imaging (MRSI) as an alternative [37]. With MRSI, a grid of spectra are acquired. MRSI has an important advantage – it provides information about the spatial distribution of metabolites, which is useful for studying multiple lesions or evaluating the spatial variation of a metabolite in a heterogeneous lesion. However, MRSI for breast is technically more challenging than SVS and quantification of metabolite levels is more problematic. For these reasons, most MRS research on breast cancer has used SVS. Figure 1 shows a representative example of a single-voxel ^1H spectrum of an invasive ductal carcinoma, with the tCho resonance indicated.




Figure 1 - Example of a localized breast ^1H spectrum of human breast acquired at 4 Tesla. The fat-suppressed, dynamic contrast-enhanced MR image (sagittal view) on the left shows the location of the VOI (red box), which covers a rim-enhancing invasive ductal carcinoma. The water-suppressed spectrum on the right shows the resonances observed in this malignant breast lesion.

While a number of groups have had success using the *detectability* of tCho to indicate malignancy, this approach assumes that the MRS measurement sensitivity is roughly constant from one measurement to the next. However there are a number of factors which make this assumption questionable in breast MRS, such as variations in voxel size, adipose tissue content, and coil sensitivity. In our experience at 4 Tesla, we have found that these factors produce a large variation in sensitivity [38]. Thus, some form of quantification should be used to correct for sensitivity variations, or at the very least exclude those voxels with unusual sensitivity.

A variety of approaches have been used for quantifying MRS data. We chose to use the intravoxel water resonance as an internal reference, because this method is robust and automatically compensates for variations in many factors [38]. Some groups have proposed using an external standard for referencing [31, 39]. This approach also works, but requires additional corrections for voxel size, adipose tissue content, and coil efficiency. Both internal and external referencing methods need correction for differences in relaxation rates which are difficult to measure in individual subjects.

A typical breast MRS study is performed immediately after acquiring dynamic contrast-enhanced (DCE) MR images. Decisions about the placement of the MRS voxel are usually based on review of the lesion morphology and the kinetics of contrast agent uptake while the patient is still in the magnet. With SVS, the placement of the voxel is of critical importance. The voxel should be placed so that it contains as much of the lesion as possible while excluding other tissues such as normal fibroglandular or adipose tissue. In studies using quantitative ^1H MRS to monitor response to treatment, the voxel size and position can be adjusted to cover the same anatomical region of the tumor and the voxel size is decreased as the tumor shrinks.

The first and most studied application for breast MRS is to distinguish benign from malignant lesions prior to biopsy. The first published paper on this topic, by Roebuck *et al.* in 1998 [31], proposed the idea that tCho could be used as a marker of malignancy. A number of papers that followed continued to use this hypothesis, but performed studies with somewhat different techniques. The overall results are quite consistent. Katz-Brull *et al.* published a combined analysis of the first five publications and reported an overall sensitivity of 83% and

specificity of 85% [40]. These results are very encouraging, especially considering that the determination of malignancy was done without considering any other diagnostic or historical information that would normally be available clinically.

Other publications describing tCho detection in breast cancers did not report diagnostic specificity and sensitivity [30, 32]. While the results using a simple detectability hypothesis are encouraging, it seems likely that there are benign pathologies that also produce detectable levels of tCho. Indeed, at 1.5 Tesla, a detectable tCho resonance has been reported in fibroadenomas [33, 34, 37, 41], tubular adenomas [31, 35], and lactating subjects [33, 36].

Two recent studies have aimed to evaluate whether MRS can improve the specificity of a diagnostic breast MR exam. Huang *et al.* appended a single-voxel MRS measurement and a single-slice T_2^* -weighted perfusion measurement to a conventional DCE-MRI exam [41]. They found that the addition of MRS increased the specificity of the exam from 62.5% to 87.5%, and the further addition of the perfusion measurement raised the specificity to 100%. Our group recently performed a retrospective blinded observer performance study with four readers and 55 subjects to determine if quantitative MRS could improve the specificity and sensitivity of a DCE-MRI exam [42]. In this study, we reported that adding quantitative MRS results to a DCE MRI exam produced improvements in the sensitivity, specificity, and accuracy for all readers, and improved the inter-observer agreement between the readers.

A second promising application of breast MRS involves predicting response to treatment. Current clinical methods such as palpation and imaging rely on changes in tumor size, which typically take several weeks before any changes are detectable. Breast MRS, in contrast, detects changes in intracellular metabolism that would occur before any gross morphological change. The first report using tCho measurements to detect treatment response in breast cancer was by Jaganathan *et al.*, who observed that the tCho resonance disappeared or became smaller in 89% of subjects undergoing chemotherapy [36]. Expanding on this observation, our group performed a study designed to determine whether changes in tCho concentration ([tCho]) could provide a biomarker of clinical response as soon as 24 hours after the first dose of doxorubicin-based chemotherapy for locally-advanced breast cancer [43]. Of the first 13 patients who successfully completed the protocol without technical problems, the change in [tCho] between baseline and 24 hours after the first dose of chemotherapy showed significant positive correlation ($R=0.79$, $p=0.001$) with the change in lesion size measured at the end of four cycles of chemotherapy. The change in [tCho] within 24 hours was significantly different between responders and nonresponders ($p=0.007$) classified using RECIST (Response Evaluation Criteria in Solid Tumors). These results suggest that the change in [tCho] within 24 hours after the first dose of the drug can serve as an early indicator for predicting clinical response to treatment for locally-advanced breast cancer.

Prostate Cancer

Unlike in breast applications, the suspicious regions in the prostate cannot be as easily localized with imaging and are typically multi-focal, necessitating the use of spectroscopic imaging instead of single voxel methods. In order to cover the entire prostate while maximizing resolution and signal-to-noise, three-dimensional ^1H MRSI is the method of choice. To localize the volume of interest while minimizing chemical shift effects, point resolved spectroscopy (PRESS) is used in combination with high bandwidth outer-volume suppression (OVS) [44]. Minimizing the effects of periprostatic lipids is accomplished with spatially and spectrally

selective RF pulses [45] or with the inclusion of spectrally selective suppression RF pulses [46-48].

Cancer in the prostate is characterized by increased tCho and decreased Cit levels. The decrease in citrate results from both a change in normal epithelial cell function and loss of ductal structure, while increased tCho levels, although not fully understood, may be due to increased cellular proliferation and density. Thus the ratio tCho/Cit is doubly sensitive to the abnormal metabolism, cellular function and morphology observed in cancer. As it is difficult in many instances to separate the resonances of tCho and tCr the ratio is typically reported as (tCho+tCr)/Cit.

Metabolic information provided by spectroscopic imaging has shown promise in targeting biopsy [49], following response to therapy [50], treatment planning [51, 52] and cancer staging [53]. Zakian et al has recommended using the combination of the maximum ratio (tCho + tCr)/Cit and the MR spectroscopic imaging tumor volume as an index to help predict tumor aggressiveness [54].




Figure 2 – T₂-weighted axial MRI (left) and selected ¹H spectra (right) from a patient with prostate cancer (Gleason score 6). The anterior part of the left peripheral zone has normal spectra (solid line) and is bright on the MR image, while the posterior par has cancerous spectra (dotted line). Reproduced from Noworolski et al, *Magn. Reson. Med.* 53:249-255 (2005).

Brain Tumors

Several early studies have suggested a role for *in vivo* ¹H MRS in helping to diagnose and monitor treatment response in brain tumors [55-60]. More recent publications have convincingly shown the value of ¹H MRSI in the clinical management of brain tumor patients in a number of areas. These include characterizing and classifying tumors [61], identifying tumor in T₂-hyperintense regions that did not display enhancement with Gd-contrast injection [62], differentiating cerebral necrosis from tumor progression [63, 64], monitoring treatment response [65-68], and predicting prognosis [69]. Similar to the prostate, 3D ¹H MRSI is generally considered to be the method of choice for evaluating brain tumors.

As compared to normal brain, common distinguishing spectral features of brain tumors include decreased levels of NAA and tCr, and elevated levels of tCho, lactate and lipids. The level of mIns, an astrocytic marker and putative osmolyte, has been reported to be elevated in hemangiopericytomas [70].




Figure 3 – Spoiled gradient-echo image (left) showing the placement of the PRESS-selected box and the MRSI grid used to obtain full coverage of the brain tumor mass. The PRESS-selected box includes the subcutaneous lipid layer at the bottom-right corner. By chopping the corner of the PRESS box using very selective OVS pulses (black line indicates edge of suppression band), contamination from mobile lipid signals arising in the scalp were avoided. Elevated tCho levels occurred in the viable tumor regions, while the absence of metabolites in other regions was indicative of necrosis. Reproduced from Tran et al, *Magn. Reson. Med.* 43:23-33 (2000).

Conclusions

The quality and reliability of MRS data will only improve as 3 Tesla systems become more common and as further refinements in techniques and software occur. Based on the convincing results obtained from multiple institutions to date, it is clear that ^1H MRS is destined to play a prominent role in the clinical management of cancer patients.

Acknowledgements

The NIH (grants CA92004, RR08079, RR00400), the MIND Institute, and the W.M. Keck Foundation are gratefully acknowledged for supporting portions of this work.

References

1. Schwarz, A.J., et al., *Early in vivo detection of metabolic response: a pilot study of ^1H MR spectroscopy in extracranial lymphoma and germ cell tumours*. Br J Cancer, 2002. **75**: p. 959-966.
2. Li, C.-W., et al., *Quantification of choline compounds in human hepatic tumors by proton MR spectroscopy at 3 T*. Magn Reson Med, 2005. **53**: p. 770-776.
3. Adalsteinsson, E., et al., *Feasibility study of lactate imaging of head and neck tumors*. NMR Biomed, 1998. **11**: p. 360-369.
4. Bourne, R.M., et al., *In vivo and ex vivo proton MR spectroscopy of primary and secondary melanoma*. Eur J Radiol, 2005. **53**: p. 506-513.
5. Mackinnon, W.B., et al., *Fine-needle biopsy specimens of benign breast lesions distinguished from invasive cancer ex vivo with proton MR spectroscopy*. Radiology, 1997. **204**: p. 661-666.
6. Cheng, L.L., et al., *Evaluating human breast ductal carcinomas with high-resolution magic-angle spinning proton magnetic resonance spectroscopy*. J Magn Reson, 1998. **135**: p. 194-202.
7. Gribbestad, I.S., et al., *Metabolite composition in breast tumors examined by proton nuclear magnetic resonance spectroscopy*. Anticancer Res, 1999. **19**: p. 1737-1746.
8. Smith, T.A., et al., *The phosphocholine and glycerophosphocholine content of an oestrogen-sensitive rat mammary tumour correlates strongly with growth rate*. Br J Cancer, 1991. **64**: p. 821-826.
9. Herminghaus, S., et al., *Increased choline levels coincide with enhanced proliferative activity of human neuroepithelial brain tumors*. NMR Biomed., 2002. **15**: p. 385-392.
10. Katz-Brull, R., et al., *Metabolic markers of breast cancer: enhanced choline metabolism and reduced choline-ether-phospholipid synthesis*. Cancer Res, 2002. **62**: p. 1966-1970.
11. de Molina, A.R., et al., *Choline kinase activation is a critical requirement for the proliferation of primary human mammary epithelial cells and breast tumor progression*. Cancer Res, 2004. **64**: p. 6732-6739.
12. Sitter, B., et al., *High-resolution magic angle spinning MRS of breast cancer tissue*. NMR Biomed, 2002. **15**: p. 327-337.
13. Barton, S.J., et al., *Comparison of in vivo ^1H MRS of human brain tumours with ^1H HR-MAS spectroscopy of intact biopsy samples in vitro*. MAGMA, 1999. **8**: p. 121-128.
14. Cheng, L.L., et al., *Non-destructive quantitation of spermine in human prostate tissue samples using HRMAS ^1H NMR spectroscopy at 9.4 T*. FEBS Lett, 2001. **494**: p. 112-116.
15. Kuesel, A.C., et al., *^1H MRS of high grade astrocytomas: Mobile lipid accumulation in necrotic tissue*. NMR Biomed, 1994. **7**: p. 149-155.
16. Ignasi, B., et al., *The relationship between nuclear magnetic resonance-visible lipids, lipid droplets, and cell proliferation in cultured C6 cells*. Cancer Res, 1999. **59**: p. 1861-1868.
17. Blankenberg, F.G., et al., *Detection of apoptotic cell death by proton nuclear magnetic resonance spectroscopy*. Blood, 1996. **87**: p. 1951-1956.
18. Hakumaki, J.M., et al., *^1H MRS detects polyunsaturated fatty acid accumulation during gene therapy of glioma: Implications for the in vivo detection of apoptosis*. Nature Med., 1999. **5**: p. 1323-1327.

19. Negendank, W.,R. Sauter, *Intratumoral lipids in ^1H MRS in vivo in brain tumors: experience of the Siemens cooperative clinical trial*. Anticancer Res, 1996. **16(3B)**: p. 1533-1538.
20. Bolan, P.J., et al., *Eliminating spurious lipid sidebands in ^1H MRS of breast lesions*. Magn Reson Med, 2002. **48**: p. 215-222.
21. Tkac, I.,R. Gruetter, *Methodology of ^1H NMR spectroscopy of the human brain at very high magnetic field*. Appl Magn Reson, 2005. **29**: p. 139-157.
22. Matsumura, A., et al., Clin Neuroncol Neurosurg, 2005. **107**: p. 379-384.
23. Terpstra, M., et al., *Relationships among lactate concentration, blood flow, and histopathologic profiles in rat C6 glioma*. NMR Biomed., 1996. **9**: p. 185-194.
24. Schwickert, G., et al., *Correlation of high lactate levels in human cervical cancer with incidence of metastasis*. Cancer Res., 1995. **55**: p. 4757-4759.
25. Walenta, S., et al., *Correlation of high lactate levels in head and neck tumors with incidence of metastasis*. Am. J. Pathol., 1997. **150**: p. 409-415.
26. Daly, P.F.,J.S. Cohen, *Magnetic resonance spectroscopy of tumors and potential in vivo clinical applications: a review*. Cancer Res, 1989. **49**: p. 770-779.
27. Leach, M.O., et al., *Measurements of human breast cancer using magnetic resonance spectroscopy: a review of clinical measurement and a report of localized ^{31}P measurements of response to treatment*. NMR Biomed, 1998. **11**: p. 314-340.
28. Sijens, P.E., et al., *Human breast cancer in vivo: H-1 and P-31 MR spectroscopy at 1.5 T*. Radiology, 1988. **169**: p. 615-620.
29. Jagannathan, N.R., et al., *Volume localized in vivo proton MR spectroscopy of breast carcinoma: variation of water-fat ratio in patients receiving chemotherapy*. NMR Biomed, 1998. **11**: p. 414-422.
30. Thomas, M.A., et al., *Volume-localized two-dimensional correlated magnetic resonance spectroscopy of human breast cancer*. J Magn Reson Imaging, 2001. **14**: p. 181-186.
31. Roebuck, J.R., et al., *Human breast lesions: Characterization with proton MR spectroscopy*. Radiology, 1998. **209**: p. 269-275.
32. Gribbestad, I.S., et al., *In vivo ^1H MRS of normal breast and breast tumors using dedicated double breast coil*. J Magn Reson Imaging, 1998. **8**: p. 1191-1197.
33. Kvistad, K.A., et al., *Characterization of neoplastic and normal human breast tissues with in vivo ^1H MR spectroscopy*. J Magn Reson Imaging, 1999. **10**: p. 159-164.
34. Yeung, D.K., et al., *Human breast lesions: characterization with contrast-enhanced in vivo proton MR spectroscopy - initial results*. Radiology, 2001. **220**: p. 40-46.
35. Cecil, K.M., et al., *The evaluation of human breast lesions with magnetic resonance imaging and proton magnetic resonance spectroscopy*. Breast Cancer Res Treat, 2001. **68**: p. 45-54.
36. Jagannathan, N.R., et al., *Evaluation of total choline from in vivo volume localized proton MR spectroscopy and its response to neoadjuvant chemotherapy in locally advanced breast cancer*. Br J Cancer, 2001. **84**: p. 1016-1022.
37. Jacobs, M.A., et al., *Proton magnetic resonance spectroscopic imaging of human breast cancer: A preliminary study*. J. Magn. Reson. Imaging, 2004. **19**: p. 68-75.
38. Bolan, P.J., et al., *In vivo quantification of choline compounds in the breast with ^1H MR spectroscopy*. Magn Reson Med, 2003. **50**: p. 1134-1143.

39. Bakken, I.J., et al., *External standard method for the in vivo quantification of choline-containing compounds in breast tumors by proton MR spectroscopy at 1.5 Tesla*. Magn Reson Med, 2001. **46**: p. 189-192.
40. Katz-Brull, R., et al., *Clinical utility of proton magnetic resonance spectroscopy in characterizing breast lesions*. J Natl Cancer Inst, 2002. **94**: p. 1197-1203.
41. Huang, W., et al., *Detection of breast malignancy: diagnostic MR protocol for improved specificity*. Radiology, 2004. **232**: p. 585-591.
42. Meisamy, S., et al., *The value of adding quantitative ¹H MRS for improving diagnostic accuracy of breast MRI: Preliminary results from an observer performance study at 4 Tesla*. Radiology, 2005. **236**: p. 465-475.
43. Meisamy, S., et al., *Predicting response to neoadjuvant chemotherapy of locally advanced breast cancer with in vivo ¹H MRS: A pilot study at 4 Tesla*. Radiology, 2004. **233**: p. 424-431.
44. Tran, T.-K.C., et al., *Very selective suppression pulses for clinical MRSI studies of brain and prostate cancer*. Magn Reson Med, 2000. **43**: p. 23-33.
45. Schricker, A.A., et al., *Dualband spectral-spatial RF pulses for prostate MR spectroscopic imaging*. Magn Reson Med, 2001. **46**: p. 1079-1087.
46. Males, R.G., et al., *Clinical application of BASING and spectral/spatial water and lipid suppression pulses for prostate cancer staging and localization by in vivo 3D ¹H magnetic resonance spectroscopic imaging*. Magn Reson Med, 2000. **43**: p. 17-22.
47. Star-Lack, J., et al., *Improved water and lipid suppression for 3D PRESS CSI using RF band selective inversion with gradient dephasing (BASING)*. Magn Reson Med, 1997. **38**: p. 311-321.
48. Futterer, J.J., et al., *Initial experience of 3 tesla endorectal coil magnetic resonance imaging and ¹H-spectroscopic imaging of the prostate*. Invest Radiol, 2004. **39**: p. 671-680.
49. Prando, A., et al., *Prostatic biopsy directed with endorectal MR spectroscopic imaging findings in patients with elevated prostate specific antigen levels and prior negative biopsy findings: early experience*. Radiology, 2005. **236**: p. 903-910.
50. Pickett, B., et al., *Time to metabolic atrophy after permanent prostate seed implantation based on magnetic resonance spectroscopic imaging*. Int J Radiat Oncol Biol Phys, 2004. **59**: p. 665-673.
51. Pouliot, J., et al., *Inverse planning for HDR prostate brachytherapy used to boost dominant intraprostatic lesions defined by magnetic resonance spectroscopy imaging*. Int J Radiat Oncol Biol Phys, 2004. **59**: p. 1196-1207.
52. Pickett, B., et al., *Static field intensity modulation to treat a dominant intra-prostatic lesion to 90 Gy compared to seven field 3-dimensional radiotherapy*. Int J Radiat Oncol Biol Phys, 1999. **44**: p. 921-929.
53. Coakley, F.V., et al., *Magnetic resonance imaging and spectroscopic imaging of prostate cancer*. J Urol, 2003. **170**: p. S69-75; discussion S75-66.
54. Zakian, K.L., et al., *Correlation of proton MR spectroscopic imaging with Gleason score based on sept-section pathologic analysis after radical prostatectomy*. Radiology, 2005. **234**: p. 804-8114.
55. Bruhn, H., et al., *Noninvasive differentiation of tumors with use of localized ¹H MR spectrscopy in vivo: initial experience in patients with cerebral tumors*. Radiol., 1989. **172**: p. 541-548.

56. Gill, S.S., et al., *Proton MR spectroscopy of intracranial tumours: in vivo and in vitro studies*. J. Comput. Assist. Tomogr., 1990. **14**(4): p. 497-504.
57. Segebarth, C.M., et al., *Detection of metabolic heterogeneity of human intracerebral tumors in vivo by ^1H NMR spectroscopic imaging*. Magn. Reson. Med., 1990. **13**: p. 62-76.
58. Kugel, H., et al., *Human brain tumors: spectral patterns detected with localized H-1 MR spectroscopy*. Radiology, 1992. **183**: p. 701-709.
59. Fulham, M.J., et al., *Mapping of brain tumor metabolites with proton MR spectroscopic imaging: clinical relevance*. Radiology, 1992. **185**: p. 675-686.
60. Ott, D., et al., *Human brain tumors: assessment with in vivo proton MR spectroscopy*. Radiology, 1993. **186**: p. 745-752.
61. Preul, M.C., et al., *Accurate, noninvasive diagnosis of human brain tumors by using proton magnetic resonance spectroscopy*. Nat Med, 1996. **2**: p. 323-325.
62. McKnight, T.R., et al., *Histopathological validation of a three-dimensional magnetic resonance spectroscopy index as a predictor of tumor presence*. J Neurosurg, 2002. **97**: p. 794-802.
63. Taylor, J.S., et al., *Clinical value of proton magnetic resonance spectroscopy for differentiating recurrent or residual brain tumor from delayed cerebral necrosis*. Int J Radiation Oncology Biol Phys, 1996. **36**: p. 1251-1261.
64. Rabinov, J.D., et al., *In vivo 3-T MR spectroscopy in the distinction of recurrent glioma versus radiation effects: Initial experience*. Radiology, 2002. **225**: p. 871-879.
65. Lazareff, J.A., et al., *Variation of post-treatment H-MRSI choline signal intensity in pediatric gliomas*. J Neuro-Oncol, 1999. **41**: p. 291-298.
66. Preul, M.C., et al., *Using proton magnetic resonance spectroscopic imaging to predict in vivo the response of recurrent malignant gliomas to tamoxifen chemotherapy*. Neurosurg, 2000. **46**: p. 306-318.
67. Kimura, T., et al., *Evaluation of the response of metastatic brain tumors to stereotactic radiosurgery by proton magnetic resonance spectroscopy, $^{201}\text{TlCl}$ single-photon emission computerized tomography, and gadolinium-enhanced magnetic resonance imaging*. J Neurosurg, 2004. **100**: p. 835-841.
68. Balmaceda, C., et al., *Multisection ^1H magnetic resonance spectroscopic imaging assessment of glioma response to chemotherapy*. J Neuro-Oncol, 2005.
69. Li, X., et al., *Identification of MRI and ^1H MRSI parameters that may predict survival for patients with malignant gliomas*. NMR Biomed, 2004. **71**: p. 10-20.
70. Barba, I., et al., *Magnetic resonance spectroscopy of brain hemangiopericytomas: high myo-inositol concentrations and discrimination from meningiomas*. J Neurosurg, 2001. **94**: p. 55-60.